

Bioarchaeological investigations at Kolonna, Aegina (Early Helladic III to Late Helladic III)

ALFRED GALIK, GERHARD FORSTENPOINTNER, GERALD E. WEISSENGRUBER, URSULA THANHEISER,
MICHAEL LINDBLOM, RUDOLFINE SMETANA & WALTER GAUß

Abstract

The interdisciplinary bioarchaeological investigations at Kolonna, Aegina yielded significant evidence of agriculture and husbandry. A clearly recognisable emphasis on ovicaprines distinguishes the livestock associated with the MH II Large Building Complex from more broadly-based or even cattle-focused economies at other contemporaneous mainland sites. A similar emphasis on small ruminants can be found in faunal samples from Aegean islands. Mycenaean layers inside and surrounding a kiln contained large numbers of heavily crushed and partly burnt shell fragments of the common purple snail, indicating most likely the process of purple dye production. The presence of few but significant remains of large game such as red deer, wild boar and lion prove hunting activities by high-ranking inhabitants of the Large Building Complex. Hunting took place most likely in the adjacent mainland regions. Game birds are represented by cormorants and pigeons, again demonstrating similarities to Middle Minoan Kommos. Seeds of ruderal plants and microfauna remains enable cautious conclusions on environmental conditions in Middle Bronze Age Aegina.

Keywords

Aegean island; Bronze Age; bioarchaeology; integrated sciences; agriculture.

Introduction

The results discussed here are based on excavations at Kolonna, Aegina mainly carried out between 2002 and 2007, at the so-called South Slope in the centre of the prehistoric settlement.¹ The Kolonna settlement on the island of Aegina has long been known as one of the major centres of the Aegean Bronze Age.² Its importance is partly due to its 'ideal' geographical position in the centre of the Saronic Gulf, at the crossroads of important Bronze Age routes between central mainland Greece, the north-east Peloponnese, the Cyclades and Crete.³ The settlement of Kolonna

flourished for almost a millennium, from the advanced Early Helladic II period (hereafter EH) through the Middle Helladic period (hereafter MH) until at least the Shaft Grave period (early Late Helladic period, hereafter LH).⁴ This is well illustrated by impressive fortifications (since the EH III period) and by a wealth of material remains.⁵ These include the earliest known shaft grave so far in the Aegean,⁶ the famous 'treasure of Aegina' now at the British Museum in London,⁷ another recently found rich metal treasure,⁸ a number of other 'prestige' items of various types, mainly of Cretan origin or influence, and substantial amounts of imported ceramic vessels from various sources across the central and southern Aegean.⁹ This evidence indicates that Aegina was an important hub in the central Aegean network, acting both as producer and, probably, distributor of commodities in a wider region. At the same time the Kolonna community was an avid consumer of imported or exotic goods while also being open to 'foreign' practices and, possibly, craftspeople. Furthermore, the first half of the second millennium saw growing indications of wealth and prestige accumulation in the hands of a certain section of the society. A MH II monumental shaft grave was built and richly furnished with weapons, jewellery

¹ For preliminary reports, see Felten *et al.* 2003, 2004, 2005, 2006, 2007, 2008, 2009.

² Rutter 1993. For a recent summary on the prehistory of Kolonna, see also Gauß 2009. For the absolute chronology see Wild *et al.* 2010.

³ See also Gauß & Kiriati in press.

⁴ See Rutter 1993 and Gauß & Kiriati in press.

⁵ Walter & Felten 1981.

⁶ Kilian-Dirlmeier 1997.

⁷ Higgins 1979, 1987; see Gauß 2006, 437 n. 2 with further references, and Phillips 2008.

⁸ Reinholdt 2008.

⁹ See similar summaries at Gauß & Kiriati in press; Gauß 2009.

and ceramic vessels,¹⁰ while the MH I-II ‘Large Building Complex’ (also referred to as ‘Mansion’) in the centre of the settlement was already in use.¹¹ The settlement was surrounded by impressive fortifications throughout the MH period. This combined evidence, implying unprecedented mobilisation of a work force and complex planning and organisation of crafts and trade activities, clearly distinguishes Kolonna from a standard MH community.¹² The Kolonna settlement seems to have been one of the earliest examples of complex society in the Aegean outside Crete, and a significant commercial and possibly political centre in the Saronic Gulf.¹³ Jeremy Rutter argued that “[...] Kolonna has emerged as a Middle Helladic site without peer on the Greek mainland”, while Wolf-Dietrich Niemeier even assumed the emergence of a ‘state’ on Aegina, comparable to contemporary political entities on Crete.¹⁴

In the context of the long exploration history of the site the focus of this new research was to integrate archaeological and bioarchaeological studies, in order to learn more about subsistence, diet and life at this important Bronze Age site. It should furthermore be stressed that the integrated archaeological and bioarchaeological analysis of material originating from a continuous stratigraphic sequence, covering the EH III to LH I periods, is thus far limited to only a few sites in mainland Greece. Therefore our analysis will hopefully help to better understand subsistence and life (including changes in diet) in these periods. Keeping in mind the statement about Kolonna being an atypical MH site, it will furthermore be very interesting to see if this status is also reflected in the recently excavated bioarchaeological remains.¹⁵

¹⁰ Kilian-Dirlmeier 1997.

¹¹ Gauß & Smetana 2010.

¹² Gauß & Kiriati in press.

¹³ Gauß & Kiriati in press.

¹⁴ Rutter 1993, 776, 780; Niemeier 1995, 74. For similar discussions, see also Dickinson 1977, 33 and Lindblom 2001, 35.

¹⁵ See for example analyses of material from Lerna in the Argolid (Gejvall 1969). See also analyses carried out at Pefkakia Maglousa (Amberger 1979), Argissa Magoula (Boessneck 1962), Ay. Mamas/Olynthos (Becker 2008) and Asine (Moberg Nilsson 1996; Nordquist 1987).

¹⁶ See Jacomet & Kreuz 1999, 62, among others.

¹⁷ For the Kolonna V destruction see Walter & Felten 1981; Gauß 2009; Wild *et al.* 2010.

¹⁸ For the find locations of barley see Walter & Felten 1981.

¹⁹ However, such pure stock with less than 1% elements other than grain is unusual. The samples were taken at the initial stages of the excavation in the late 1960s and 1970s, when sampling for archaeobotany was no common procedure, and they may have been sieved with a coarse mesh with a mesh size larger than 2mm.

Precise excavation techniques were applied during the 2003-2007 excavations at Kolonna including flotation of ca. 5m³ of soil on a 0.5-mm screen, and the sieve residues revealed huge amounts of bioarchaeological remains. Most of the material discussed here was associated with the Large Building Complex, presumably the mansion of the prehistoric settlement, and dates to MH II. The research also focused on bioarchaeological remains of the EH III and the LH III period. The aim of all these investigations was to answer economical and ecological questions from an integrated bioarchaeological point of view.

Plant remains

Agriculture certainly made an important contribution to the nutrition of the inhabitants of the island. In contrast to animal bone and molluscs, however, vegetal food remains are scarce. They mainly derive from dated layers (fills) but sometimes cannot be associated with specific contexts.

Plant remains had been preserved in two forms, mineralised and charred. In an environment rich in calcium carbonate and phosphate, as may be found in sewers or latrines, deposited plant parts will become impregnated with these carbonates and phosphates. After the decay of the organic components, the plants are present as mineral, glass-like casts.¹⁶ In Kolonna only fig (*Ficus carica*) and grape (*Vitis vinifera*) are present in mineralised form, with fig pips dominating. They are abundant throughout the Bronze Age site but are most numerous in areas with a low concentration of charred plant remains. Charring is the result of a burning event by which the organic material is reduced to almost pure carbon. Charred plant remains in domestic contexts can usually be linked to food preparation, the use of plants as fuel or the accidental burning of structures. An unusual find comes from the late EH III phase, where a high amount of charred barley (*Hordeum vulgare*) was recovered in the burnt houses of the Kolonna V settlement.¹⁷ The assemblage consisted of about three litres of grain and some detached glumes, while rachis segments and weeds were absent. In another location within the same destruction area about two litres of fine grey ash were recovered, and phytolith analysis revealed that cereal straw got burnt there.¹⁸ The area seems to have served as a storage facility where highly processed and thoroughly cleaned barley, ready for consumption, plus straw were kept in separate places awaiting further use.¹⁹ In all other areas

of the site the density of plant remains is very low. The average 10-litre soil sample from the Large Building Complex, for example, would yield 17 items in MH II deposits and six items in MH III strata. In both periods mineralised fig pips were dominant. Wheat (einkorn – *Triticum monococcum* and emmer – *T. dicoccum*) and barley were found in equal proportions. Pulses are represented by lentils (*Lens culinaris*) and peas (*Pisum sativum*), and also by the slightly poisonous bitter vetch (*Vicia ervilia*) and other unidentifiable legume species. Fig and olive trees – the latter rarely attested – and grape vines provided fruit. Although scarce, the plant remains indicate that the plant food economy was broad-based, with a variety of cereals and pulses providing carbohydrates and protein,²⁰ and fruits and nuts adding variety to the diet. The finds may point to a subsistence economy based on agricultural production,²¹ with parallels on mainland Greece.

Mammal and bird remains

The animal protein sources in the MH II Large Building Complex resembled those at comparable island or mainland coastal sites. They were varied but seem to have been dominated by the exploitation of the major domesticates. Numerical quantification of the major domesticates revealed an emphasis on ovicaprines, also in contexts other than the MH II Large Building Complex, with a clear preference for goats (Figures 1 and 2).²² Even the weight distribution underlines the importance of ovicaprines at Kolonna. It displays rather high values compared to the weight distribution of cattle remains. The exploitation of pigs was certainly important but to a lesser extent. The distribution of represented body parts of the major domesticates reveals a rather unspecific use of the complete carcass of the major domesticates at the settlement (Figures 1 and 2).

Pigs were mainly slaughtered as juveniles and premature individuals, and only a few remains suggest the culling of piglets. The pattern of ovicaprines with regard to the culling of kids and lambs appears similar, but the number of premature to adult slaughtered individuals is significantly higher. As pigs were predominantly kept for meat production, the culling profile of ovicaprines allows the possibility that they were used for renewable products like dairy, hair and wool.

Cattle were predominantly culled at premature (before the late fusion of epiphyses, i.e. prior to 3.5 to 4 years) and mature ages (older than 4 years), indicating




FIGURE 1. Numerical quantification of major domesticates associated with the MH II Large Building Complex.




FIGURE 2. Quantification of major domesticates associated with the MH II Large Building Complex by bone weight.

the use of the animals’ work power, although only one proximal phalanx (associated with the MH II Large Building Complex) shows proximo-cranial lipping at a medium stage and widened palmar ligament bases.

²⁰ As in other mainland sites, such as Tiryns (Kroll 1982), Asine (Nordquist 1987) or Olynthos (Kroll 2008).

²¹ Renfrew 1972, 282-284.

²² Gamble 1978, 749-750 reports more horn cores of goat than those of sheep from MM (Middle Minoan) Thera. The MBA (Middle Bronze Age) assemblage from mainland Olynthos indicates a rather different behaviour, favouring sheep (Becker 2008, 171). Some literature on distinguishing sheep from goat: post-cranial – Boessneck 1969; Boessneck *et al.* 1964; Buitenhuis 1995; Kratochvil 1969; Payne 1969; Prummel & Frisch 1986; Zeder & Lapham 2010; dentition – Brouwer 2002; Halstead & Collins 2002; Zeder & Pilaar 2010.

Exploitation and herding strategies at MH Greek mainland sites involved a more balanced use of the major domesticates,²³ while the pattern observed at the MH II Large Building Complex in Kolonna suggests live stock management similar to that on Minoan Crete. In Middle Minoan II (hereafter MM) Kommos the proportion of ovicaprines was over 60%. Reese lists several sites at Minoan Crete which show the same husbandry pattern for the three major domesticates as that at Kommos or Aegina.²⁴ Keeping small ruminants on small islands is certainly less problematic than to keep and feed cattle herds.²⁵

A shallow pit associated with the MH II Large Building Complex contained a more or less completely preserved ram skeleton. Heavy chopping marks on the neck vertebrae probably indicate how the animal was killed. After that, the chest of this individual was carefully opened, leaving butchering marks on the sternum. This might indicate ritual

activities that involved the sacrificial killing of live stock.

The thick infill layer associated with the MH II Large Building Complex contained mainly domestic refuse and yielded various remains of big game. The most outstanding find is undoubtedly a lion humerus, that suggests a rather large male individual (proximal width = 73mm, proximal thickness = 85mm). This find places Aegina Kolonna amongst the six widespread locations from the Late Neolithic to the Archaic Age that testify to the presence of lions in Greece.²⁶ The bone does not show any traces of dismembering, but the fracture edges reveal perimortal smashing as they still contained collagen. The proximal section certainly suggests a meat bearing part, rather than a fragment that was part of an imported trophy. Morphological analysis enabled the taxonomical assignment of the upper arm fragment to the Asian Lion (*Panthera leo persica*), which was also the case at Aegira,²⁷ Kalapodi,²⁸ and Kastanas.²⁹ Remains of big game associated with the MH II Large Building Complex are generally scarce. However, the postcranial bones indicate consumption of prestigious animals like red deer (*Cervus elaphus*), fallow deer (*Dama dama*), roe deer (*Capreolus capreolus*), wild boar (*Sus scrofa*) and most probably also aurochs (*Bos primigenius*).³⁰ While the existence of native wild boar and probably roe deer as well as fallow deer on MBA Aegina is not implausible, the ecological conditions on such a small island are definitely not advantageous for the survival of a herd of stags or a pack of lions.

Besides hunting big game the Kolonna people, and presumably especially the social elite, hunted game birds as well. A few skeletal remains associated with the Large Building Complex indicate the presence of rock partridge (*Alectoris graeca*), little owl (*Athene noctua*) and greylag goose (*Anser anser*), birds that occur in Greek mainland assemblages as well as island sites. The MH II infill of the Large Building Complex yielded pigeon (*Columba* sp.) (n=8, MNI=4) and cormorant (*Phalacrocorax* sp.) bones (n=3, MNI=2), identified by butchering marks as food remains.³¹ Bones of pigeon and shearwater dominate the avian assemblage at MM Kommos.³² Aquatic birds like cormorants or even shearwaters are mainly fish eaters and their flesh therefore tastes fishy, indicate possibly an appreciation of wildfowl dishes, in particular those containing greasy and fishy bird species such as those attested at Kommos and Aegina Kolonna.

²³ Macedonia: Kastanas (Becker 1986), Olynthos (Becker 2008), Sitagroi (Bökönyi 1986). Thessaly: Argissa Magula (Boessneck 1962), Magula Pevkakia (Jordan 1975; Amberger 1979). Peloponnese: Tiryns (Van den Driesch & Boessneck 1990), Lerna (Gejvall 1961, Tab. 3), Nichoria (Sloan & Duncan 1978), Asine (Moberg Nilsson 1996, 400).

²⁴ Crete: Reese *et al.* 1995, 166. Reese compares various sites on Minoan Crete and reaches the same conclusion, i.e. that the islanders based their mammalian protein intake on ovicaprine stock. Similar patterns are reported for the Cyclades: Phylakopi (Gamble 1985, 169), Akrotiri (Gamble 1978, 750; Trantalidou 1990, 400).

²⁵ As discussed by Gamble 1978, 150.

²⁶ Evidence such as loose teeth from Pylos, Ayia Irini or Mycenae, or distal autopodial bones from Samos and Delphi were most likely imported as trophies or amulets (Thomas 2004, 190), and do not necessarily indicate the presence of lions near these sites. However, lion bones of 'meat bearing' parts like the humerus from Aegina may prove the presence of these animals in the vicinity of sites, such as at BA to Early Archaic Tiryns (Van den Driesch & Boessneck 1990, 110), Kastanas (Becker 1986, 336), Hyampolis-Kalapodi (Van den Driesch & Boessneck 1990, 110), Late Neolithic Dikili Tash (Helmer 1997, 41) and Aigira (Forstenpointner *et al.* 2010, 180-181).

²⁷ For a detailed description of the Asian Lion remains, see Forstenpointner *et al.* 2010, 180-181.

²⁸ Stanzel 1991.

²⁹ Becker 1986.

³⁰ Trantalidou 2000, 718-721 provides an overview showing that red deer and fallow deer as well as roe deer and wild boar were the usual game animals at various prehistoric Greek sites. Evidence for urochs is reported from Olynthos (Becker 2008, 87), Magoula Pevkakia (Amberger 1969), Lerna (Gejvall 1969) and Argissa Magoula (Boessneck 1962).

³¹ Becker 2008, 103 presents more evidence for cormorant from Olynthos.

³² Reese *et al.* 1995, 195, Tab. 5.9.

Shells

The frequency of shell finds reflects the importance of molluscs as a protein source in Bronze Age Aegina.³³ However, the majority of the terrestrial gastropods (*Eobania vermiculata*, *Helicella* sp., *Rumina decollata*) represent secondary intrusions of modern species. Even most of the shells of the large garden snail (*Helix* sp.) entered the archaeological contexts secondarily. A small number of *Helix* shells appeared as a regular part of the archaeological assemblages, but this does not necessarily imply they were being exploited. In general, land snails occur only in small numbers on Greek sites.³⁴




FIGURE 3. Numerical quantification of the mollusc remains from the infill of the MH II Large Building Complex.

At the MH II Large Building Complex numerous gastropods as well as bivalves and a few cuttlefish remains reflect exploitation of marine resources (Figure 3). This combination of species is largely similar to that of the assemblages at MM Kommos,³⁵ and it is not unusual at mainland coastal sites, either.³⁶ A few generations earlier the Aeginetans of the EH III period had collected primarily limpets (mainly *Patella caerulea*) and top shells (*Monodonta turbinanta* besides a few *M. articulata* and *Gibbula* sp.; Figure 4). From the MH onwards purple snails (mainly *Hexaplex trunculus* and to a lesser extent *Bolinus brandaris*) were constantly being exploited in larger quantities (Figure 4). Other gastropods like tun snail (*Tonna galea*), triton (*Charonia* sp.) and ceriths (*Gourmya vulgata*) also suggest consumption.

Spondylus (*Spondylus gaederopus*), Noah’s Arc shell (*Arca noae*) and few specimens of bearded Arc shell




FIGURE 4. Quantification of gastropod remains by chronological subunit from trenches Q3 and Q6. The values in brackets indicate the estimated Minimum Number of Individuals (MNI).

(*Barbatia barbata*) were regularly used as food in the MH II Large Building Complex and in other Bronze Age contexts at Aegina. The numerous remains of pen shell (*Pinna nobilis*) are certainly overrepresented due to the high degree of fragmentation of their large and fragile valves, but it may also indicate intense exploitation (Figure 5). Less frequent are other species




FIGURE 5. Numeric quantification of bivalve remains from trenches Q3 and Q6, by general chronological period. The values in brackets indicate the estimated Minimum Number of Individuals (MNI).

³³ As introduced, with preliminary results, in Galik 2010.
³⁴ Reese *et al.* 1995; Reese 1998; Karali 1999, 2005.
³⁵ Reese *et al.* 1995, 240, 273.
³⁶ Becker 2008, 110-112; Magoula Pavlina; Prummel 2003, 57-59; Theodoropoulou 2007, 74-79.

such as mussel (*Mytilus galloprovincialis*), scallop (*Pecten jacobaeus* and *Chlamys* sp.), and species preferring soft grounds such as lagoon cockle (*Cerastoderma glaucum*) and a few spiny cockles (*Acanthocardia tuberculata*) as well as Venus shells (*Venus verrucosa* and *Chamelea gallina*).

In general only a few worked shells occur amongst Bronze Age mollusc material. The abundance of pen shell may suggest the use of valves as raw material. Although spondylus was a consumed shellfish and large stocks must have been available, the Bronze Age people in Aegina did not use its shell as raw material. Some modified bivalve shells of *Glycimeris* point towards their decorative use.³⁷ Rare gastropod species like dog whelk (*Buccinulum corneum*), abalone (*Haliotis* sp.), cone shell (*Conus mediterraneus*) and cowry (*Luria lurida*) were probably also used to produce ornaments.

An abundance of remains of purple snail shell usually indicates purple dye production. The preparation to extract the secretions of the hypobranchial gland involves intense crushing of the shells, according to ancient authors³⁸ and modern experimental archaeological research.³⁹ The area around and inside a LH IIIA kiln in trench Q₃ contained much higher quantities of such remains (Figure 4).⁴⁰ Some of the fragments were burnt and probably represent a method to get rid of this kind of waste by burning it to produce lime.⁴¹ However, the question remains to what extent purple dye was produced, since the Minimum Number of Individuals (MNI) in LH IIIA is only 370 specimens. Such a relatively small number of individuals may indicate that purple dye production was limited to the scale of individual (small) households.⁴² However, the area around the LH kiln provided only limited information and the bulk of the refuse probably accumulated

somewhere else. However, even in the case of household production, several households in combination would produce a substantial amount of dye, which may explain how the economy and wealth accumulation developed in Kolonna in the Bronze Age. In a wider context, MM and MH purple dye production has been recorded at Palaikastro, Kouphonisi, Ayios Mamas, Akrotiri, Asine, Aghios Kosmas and Olynthos as well as in the entries on purple production on the Linear B tablets from Knossos.⁴³ The EH pattern of mollusc exploitation at Kolonna differs from that indicated by the MH and LH assemblages, with increasing exploitation of purple snails beginning in the MH period, and purple dye production may also have occurred in the LH period (Figure 5).

Conclusions

Kolonna gained in importance during the MH period and became the wealthy main settlement of the island. Evidence for Cycladic connections appears around EH III in the form of imported goods. Minoan influence is detectable from the MH onwards as the appearance of imported technology and goods.⁴⁴ On the other hand, exported Aeginetan pottery turns up in many central Aegean sites in the MBA. The plant remains reflect a wide range of cereals, pulses as well as fruits like figs, vine and olives, indicating a basic agricultural economy from the EBA onwards, which was comparable to that at coastal mainland sites. The inhabitants of the Large Building Complex certainly formed some kind of elite in Kolonna, which the remains of large game may confirm.⁴⁵ For these people, Minoan influence expressed itself as exploitation of economically useful animals. Contrary to a preference for cattle on the mainland, meat production at Kolonna was based more on ovicaprine exploitation like at MM Kommos, and the inhabitants of the Large Building Complex clearly shared the surprising taste for bird meat of the Cretan people.⁴⁶ The exploitation pattern of molluscs may provide more insight into chronological changes, since the composition of mollusc assemblages alters at the transition of the EH to the MH. An increase in heavily smashed purple snail shells in the LH probably points to the production of purple dye. The archaeological as well as the bioarchaeological finds from the recent excavations at Kolonna point to an intense exchange between Minoan Crete and the Cyclades. Kolonna played undoubtedly a significant role in these well-established

³⁷ Reese *et al.* 1995, 240 mentions that most of the *Glycimeris* shells were collected when the animals lay dead on the beach, which is similar to what is indicated by the assemblage from Aegina. See for Aegean shellfish modification in the Aegean Reese 1983, 1984; Karali 1990; Karali-Yannacopoulos 1992; Becker 2008.

³⁸ For example Pliny the Elder (Hist. Nat. IX), Aristoteles (Hist. an. V, 547a) or Vitruvius.

³⁹ Ruscillo 2005.

⁴⁰ For the kiln, see Gauß 2007.

⁴¹ Reese 1987; Reese *et al.* 1995, 206.

⁴² Becker 2008, 174-175 describes similar behaviour in MH Olynthos including a possible Mycenaean influence.

⁴³ Reese 1987; Reese *et al.* 1995; Karali 1999, 2008; Becker 2008.

⁴⁴ Gauß & Smetana 2010, 170.

⁴⁵ As discussed by Forstenpointner *et al.* 2010, 739.

⁴⁶ Forstenpointner *et al.* 2010, 740.

trading and exchange networks, whereby the Aeginetans used their position to trade goods – and probably to control that trade – between the islands and the coastal mainland.⁴⁷

A. GALIK

Institute for Anatomy and Histology,
University of Veterinary Medicine, Vienna
Alfred.Galik@vu-wien.ac.at

G. FORSTENPOINTNER

Institute for Anatomy and Histology,
University of Veterinary Medicine, Vienna
gerhard.forstenpointner@vu-wien.ac.at

G.E. WEISSENGRUBER

Institute for Anatomy and Histology,
University of Veterinary Medicine, Vienna
Gerald.Weissengruber@vetmeduni.ac.at

U. THANHEISER

VIAS, Vienna Institute for Archaeological Science,
University of Vienna
ursula.thanheiser@univie.ac.at

M. LINDBLOM

Department of Archaeology and Ancient History,
Uppsala University
Michael.lindblom@antiken.uu.se

R. SMETANA

Fachbereich Altertumswissenschaften /
Klassische und Frühägäische Archäologie,
University of Salzburg
rudolfine.smetana@sbg.ac.at

W. GAUß

Austrian Archaeological Institute, Athens Office
wgauss@otenet.gr

Acknowledgements

The projects were kindly funded by grants from INSTAP during the years 2003-2005 and 2007. The ready cooperation of the Institute of University of Salzburg and the Institute of Anatomy and Histology at the Veterinary Medicine University of Vienna facilitated not only a better exchange of ideas but also improved understanding between these closely linked branches of archaeological science.

References

- ALRAM-STERN, E., G. NIGHTINGALE & A.E. BÄCHLE (eds) 2007. *Keimelion. Elitenbildung und elitärer Konsum von der mykenischen Palastzeit bis zur Homerischen Epoche*. Veröffentlichungen der Mykenischen Kommission 27. Vienna.
- AMBERGER, K.-P. 1979. *Neue Tierknochenfunde aus der Magula Pevkakia in Thessalien. II. Die Wiederkäufer*. Diss. Universität München.
- BECKER C. 2008. Die archäozoologischen Befunde. In: C. Becker & H. Kroll (eds), *Das Prähistorische Olymb. Ausgrabungen in der Toumba Agios Mamas 1994-1996. Ernährung und Rohstoffnutzung im Wandel*. Prähistorische Archäologie in Südosteuropa 22, 49-184. Berlin.
- BECKER, C. 1986. *Kastanas. Ausgrabungen in einem Siedlungshügel der Bronze- und Eisenzeit Makedoniens 1975-1979. Die Tierknochenfunde*. Prähistorische Archäologie in Südosteuropa 5. Berlin.
- BOESSNECK J. 1969. Osteological Differences Between Sheep (*Ovis aries* L.) and Goat (*Capra hircus* L.). In: D. Brothwell & E. Higgs (eds), *Science in Archaeology*, 331-358. New York.
- BOESSNECK, J. 1962. Die Tierreste aus der Argissa-Magula vom präkeramischen Neolithikum bis zur mittleren Bronzezeit. In: V. Milošević, J. Boessneck & M. Hopf (eds), *Die Deutschen Ausgrabungen in der Argissa-Magula in Thessalien. I. Das präkeramische Neolithikum sowie die Tier- und Pflanzenreste*, 27-99. Bonn.
- BOESSNECK, J., H.H. MÜLLER & M. TEICHERT 1964. Osteologische Unterscheidungsmerkmale zwischen Schaf (*Ovis aries* L.) und Ziege (*Capra hircus* L.). *Kühn-Archiv* 78, 1-129.
- BÖKÖNYI, S. 1986. Faunal remains. In: C. Renfrew, M. Gimbutas & E.S. Elster, *Excavations at Sitagroi. A Prehistoric Village in Northeast Greece*, vol. 1. Monumenta Archaeologica. Los Angeles.
- BROUWER, M. 2002. *Sorting the Mandibular Teeth of Sheep from Goat. The Identification of Sheep and Goat on the Basis of Neolithic Dental Material from Eastern Syria*. MSc Diss. University of Groningen.
- BUITENHUIS, H. 1995. A Quantitative Approach to Species Determination of Ovicapridae. In: H. Buitenhuis & H.P. Uerpmann (eds), *Archaeozoology of the Near East II*, 144-150. Leiden.
- CLINE, E.H. (ed.) 2009. *The Oxford Handbook of the Bronze Age Aegean*. Oxford.
- CZERNY, E., I. HEIN, H. HUNGER, D. MELMAN & A. SCHWAB (eds) 2006. Timelines. Studies in Honour of Manfred Bietak. *Orientalia Lovaniensia Analecta* 149. Leuven, Paris, Dudley.
- DICKINSON, O.T. 1977. The Origins of Mycenaean Civilization. *Studies in Mediterranean Archaeology* 49. Göteborg.
- DRIESCH, A. v. D. & J. BOESSNECK 1990. Die Tierreste von der mykenischen Burg Tiryns bei Nauplion/Peloponnes. *Forschungen und Berichte* II, 87-164. Mainz.

⁴⁷ Philippa-Touchais *et al.* 2010.

- FELTEN, F. 2006. Ägina-Kolonna 2005. Vorbericht über die Grabungen des Fachbereichs Altertumswissenschaften / Klassische und Frühägäische Archäologie der Universität Salzburg Instituts für Klassische Archäologie der Universität Salzburg. *Jahreshefte des Österreichischen archäologischen Instituts in Wien* 75, 9-38.
- FELTEN, F. 2007. Ägina-Kolonna 2006. Vorbericht über die Grabungen des Fachbereichs Altertumswissenschaften / Klassische und Frühägäische Archäologie der Universität Salzburg Instituts für Klassische Archäologie der Universität Salzburg. *Jahreshefte des Österreichischen archäologischen Instituts in Wien* 76, 89-119.
- FELTEN, F. 2008. Ägina-Kolonna 2007. Vorbericht über die Grabungen des Fachbereichs Altertumswissenschaften / Klassische und Frühägäische Archäologie der Universität Salzburg. *Jahreshefte des Österreichischen archäologischen Instituts in Wien* 77, 47-76.
- FELTEN, F. 2009. Ägina-Kolonna 2008. Vorbericht über die Grabungen des Fachbereichs Altertumswissenschaften / Klassische und Frühägäische Archäologie der Universität Salzburg Instituts für Klassische Archäologie der Universität Salzburg. *Jahreshefte des Österreichischen archäologischen Instituts in Wien* 78, 47-76.
- FELTEN, F., S. HILLER, C. REINHOLDT, W. GAUß & R. SMETANA. 2003. Ägina-Kolonna 2002. Vorbericht über die Grabungen des Instituts für Klassische Archäologie der Universität Salzburg. *Jahreshefte des Österreichischen archäologischen Instituts in Wien* 72, 41-65.
- FELTEN, F., C. REINHOLDT, E. POLLHAMMER, W. GAUß & R. SMETANA. 2005. Ägina-Kolonna 2004. Vorbericht über die Grabungen des Fachbereichs Altertumswissenschaften / Klassische und Frühägäische Archäologie der Universität Salzburg Instituts für Klassische Archäologie der Universität Salzburg. *Jahreshefte des Österreichischen archäologischen Instituts in Wien* 74, 7-37.
- FELTEN, F., S. HILLER, C. REINHOLDT, W. GAUß & R. SMETANA. 2004. Ägina-Kolonna 2003. Vorbericht über die Grabungen des Instituts für Klassische Archäologie der Universität Salzburg. *Jahreshefte des Österreichischen archäologischen Instituts in Wien* 73, 97-128.
- FORSTENPOINTNER, G., A. GALIK, G.E. WEISSENGRUBER, S. ZOHMANN, U. THANHEISER & W. GAUß 2010. Subsistence and More in Middle Bronze Age Aegina Kolonna: Patterns of Husbandry, Hunting and Agriculture. In: A. Philippa-Touchais, G. Touchais, S. Voutsaki & J. Wright (eds), *MESOHELLADIKA: The Greek Mainland in the Middle Bronze Age*, 733-742. Bulletin de Correspondance Hellénique, Supplément LII. Athens.
- FORSTENPOINTNER, G., E. PUCHER, G.E. WEISSENGRUBER & A. GALIK 2006. Die archäozoologischen und archäobotanischen Hinterlassenschaften. In: Alram-Stern, E. & S. Deger-Jalkotzy (eds), *Aigeira I. Die mykenische Akropolis*, fascicle 3, 171-188. Österreichische Akademie der Wissenschaften, Philosophisch-Historisch Klasse, Denkschriften 342. Vienna.
- GALIK, A., S. ZOHMANN, G. FORSTENPOINTNER, G.E. WEISSENGRUBER & W. GAUß 2010. Subsistence and More in Middle Bronze Age Aegina Kolonna: Exploitation of Marine Resources. In: A. Philippa-Touchais, G. Touchais, S. Voutsaki & J. Wright J. (eds), *MESOHELLADIKA: The Greek Mainland in the Middle Bronze Age*, 743-751. Bulletin de Correspondance Hellénique, Supplément LII. Athens.
- GAMBLE, C. 1978. The Bronze Age Animal Economy from Akrotiri: a Preliminary Analysis. In: C. Doumas (ed.), *Thera and the Aegean World I*, 745-753. London.
- GAMBLE, C. 1985. Formation Processes and the Animal Bones from the Sanctuary at Phylakopi. In: C. Renfrew (ed.), *The Archaeology of Cult. The Sanctuary at Phylakopi* 479-483. The Annual of the British School at Athens Supplement vol. 18.
- GAUß, W. 2006. Minos auf Ägina. Beobachtungen zu den Beziehungen Äginas zu Kreta. In: E. Czerny, I. Hein, H. Hunger, D. Melman & A. Schwab (eds), *Timelines. Studies in honour of Manfred Bietak*. Orientalia Lovaniensia Analecta 149, 435-446. Leuven, Paris, Dudley.
- GAUß, W. 2007. Ägina Kolonna in frühmykenischer Zeit. In: Alram-Stern, E., G. Nightingale & A.E. Bächle (eds), 2007. *Keimelion. Elitenbildung und elitärer Konsum von der mykenischen Palastzeit bis zur Homerischen Epoche*, 163-172. Veröffentlichungen der Mykenischen Kommission 27. Vienna.
- GAUß, W. 2009. Aegina. In: Cline (ed.) *The Oxford Handbook of the Bronze Age Aegean*, 737-751. Oxford.
- GAUß, W. & E. KIRIATZI In press. *Pottery Production and Importation in Bronze Age Kolonna, Aegina, Greece (EBA III-LBA). An Integrated Archaeological and Scientific Study of an Island ceramic Landscape*. Ägina-Kolonna Forschungen und Ergebnisse 5. Vienna.
- GAUß, W. & R. SMETANA 2010. Aegina-Kolonna in the Middle Bronze Age. In: Philippa-Touchais, A., G. Touchais, S. Voutsaki & J. Wright (eds), *MESOHELLADIKA: The Greek Mainland in the Middle Bronze Age*, 165-174. Bulletin de Correspondance Hellénique, Supplément LII. Athens.
- GEJVALL, N.-G. 1969. *Lerna. A Preclassical Site in the Argolid. Vol. I. The Fauna*. Princeton.
- HALSTEAD P. & P. COLLINS 2002. Sorting Sheep from Goats: Morphological Distinctions Between the Mandibles and Mandibular Teeth of Adult Ovis and Capra. *Journal of Archaeological Science* 2002, 545-553.
- HELMER D. 1997. Faune sauvage et domestique: Dikili Tash à l'Époque Néolithique. *Les dossiers d'archéologie* 222, 40-41.
- HIGGINS, R.A. 1979. *The Aegina Treasure. An Archaeological Mystery*. London.
- HIGGINS, R.A. 1987. A Gold Diadem from Aegina. *Journal of Hellenic Studies* 107, 182.
- JACOMET, S. & A. Kreuz 1999. *Archäobotanik*. Stuttgart.
- JORDAN, B. 1975. *Tierknochenfunde aus der Magula Pevkakia in Thessalien*. Diss. Universität München.
- KARALI, L. 2005. Shells from the Prehistoric Sites in Northern Greece. In: D. Bar-Yosef Mayer (ed.), *Archaeomalacology: Molluscs in Former Environments of Human Behaviour*, 91-98, Oxford.
- KARALI, L. 2008. Archaeomalacological Research in Greece. <http://environmentalarchaeology.wordpress.com/2008/11/02/archaeomalacological-research-in-greece>, visited 2 March 2011.

- KARALI, L. 1999. *Shells in Aegean Prehistory*. British Archaeological Reports – International Series 761. Oxford.
- KARALI-YANNACOPOULOS, L. 1990. Sea Shells, Land Snails and Other Marine Remains from Akrotiri. In: D.A. Hardy (ed.), *Thera and the Aegean World III: Earth Sciences*, 410-415. London.
- KILIAN-DIRLMEIER, I. 1997. *Das mittelbronzezeitliche Schachtgrab von Ägina*. Alt-Ägina IV.3. Mainz-am-Rhein.
- KRATOCHVIL, Z. 1969. Species Criteria on the Distal Section of the Tibia in *Ovis ammon* F. aries L. and *Capra aegagrus* F. hircus L. *Acta Veterinaria* 38, 483-490.
- KROLL, H. 1982. Die Kulturpflanzen von Tiryns. *Archäologischer Anzeiger* 467-485.
- KROLL, H. 2008. Die archäobotanischen Befunde. In: C. Becker & H. Kroll (eds), *Das Prähistorische Olynth. Ausgrabungen in der Toumba Agios Mamas 1994-1996. Ernährung und Rohstoffnutzung im Wandel*. Prähistorische Archäologie in Südosteuropa 22, 17-48. Rahden.
- LINDBLOM, M. 2001. *Marks and Makers. Appearance, Distribution and Function of Middle and Late Helladic Manufacturers' Marks on Aeginetan pottery*. Studies in Mediterranean Archaeology 128. Jonsöred.
- MOBERG NILSSON, K. 1996. Animal Bones from Terrace III in the Lower Town of Asine. In: R. Hägg, G.C. Nordquist & B. Wells (eds), *Asine III. Supplementary Studies on the Swedish Excavations 1922-1930*, 111-115. Jonsöred.
- NIEMEIER, W.-D. (1995). Aegina. First Aegean State Outside of Crete? In: Laffineur, R. & W.-D. Niemeier (eds) 1995. *Politeia: Society and State in the Aegean Bronze Age*, 73-79. Aegaeum 12. Liège, Austin.
- NORDQUIST G.C. 1987. *A Middle Helladic Village. Asine in the Argolid*. Boreas 16.
- PAYNE, S. 1969. A Metrical Distinction Between Sheep and Goat Metacarpals. In: P.J. Ucko & G.W. Dimbleby (eds), *The Domestication and Exploitation of Plants and Animals*, 295-306. London.
- PHILLIPS, J. 2008. *Aegyptiaca on the Island of Crete in their Chronological Context. A Critical Review*. Contributions to the Chronology of the Eastern Mediterranean 18. Vienna.
- PHILIPPA-TOUCHAIS, A., G. TOUCHAIS, S. VOUTSAKI & J. WRIGHT (eds), *MESOHELLADIKA: The Greek Mainland in the Middle Bronze Age*. Bulletin de Correspondance Hellénique 508, Supplément LII. Athens.
- PRUMMEL, W. & H.J. FRISCH 1986. A Guide for the Distinction of Species, Sex, and Body Size of Sheep and Goat. *Journal of Archaeological Science* 13, 567-577.
- PRUMMEL, W. 2003. Middle Bronze Age and Hellenistic Mollusk Meals from Sourpi Bay (Thessaly, Greece). *Buletin of the Florida Museum of Natural History* 44(1), 55-64.
- REESE, D.S. 1998. Appendix II: The Faunal Remains. In: G. Walberg (ed.), *The Excavations on the Acropolis of Midea* vol. I, 1, 278-280 & 290-292. Stockholm.
- REESE, S.D. 1983. The Use of Cone Shells in Neolithic and Bronze Age Greece. *The Annual of the British School at Athens* 79, 237-238.
- REESE, S.D. 1984. Topshell Rings in the Aegean Bronze Age. *The Annual of the British School at Athens* 76, 354-357.
- REESE, D.S., M.J. ROSE & S. PAYNE 1995. The Minoan Fauna. In: J.W. Shaw & M.C. Shaw (eds), *Kommos I. The Kommos Region and Houses of the Minoan Town*, 628. Princeton.
- REESE, D. 1987. Palaikastro Shells and Bronze Age, Purple-dye Production in the Mediterranean Basin. *The Annual of the British School at Athens* 82, 201-206.
- REINHOLDT, C. 2008. *Ägina-Kolonna. Forschungen und Ergebnisse 2: Der frühbronzezeitliche Schmuckhortfund von Kap Kolonna. Ägina und die Ägäis im Goldzeitalter des 3. Jahrtausends v. Chr.* Contributions to the Chronology of the Eastern Mediterranean 15. Vienna.
- RENFREW, A.C. 1972. *The Emergence of Civilization: The Cyclades and the Aegean in the Third millenium B.C.* London.
- RUSCILLO, D. 2005. Reconstructing Murex Royal Purple and Biblical Blue in the Aegean. In: D. Bar-Yosef Mayer (ed.), *Archaeomalacology: Molluscs in former Environments of Human Behaviour*, 99-106. Oxford.
- RUTTER, J.B. 1993. Review of Aegean Prehistory, 2. The Pre-palatial Bronze Age of the Southern and Central Greek Mainland. *American Journal of Archaeology* 97, 745-797.
- STANZEL, M. 1991. *Die Tierreste aus dem Artemis-/Apollon-Heiligtum bei Kalapodi in Bötien/Griechenland*. Diss. Universität München.
- THEODOROPOULOU, T. 2007. La mer dans l'assiette: l'exploitation des faunes aquatiques dans l'alimentation en Égée Pré- et Protohistorique. In: C. Mee & J. Renard (eds), *Cooking up the Past. Food and Culinary Practices in the Neolithic and Bronze Age Aegean*, 71-88. Oxford.
- THOMAS, R.N. 2004. The Early Mycenaean Lion Up to Date. In: A.P. Chapin (ed.), *Essays in Honour of Sara A. Immerwahr*. Hesperia Suppl. 33, 161-206.
- TRANTALIDOU, K. 1990. Animals and Human Diet in the Prehistoric Aegean. In: D.A. Hardy (ed.), *Thera and the Aegean World III, vol. 2: Earth Sciences*, 392-405. London.
- TRANTALIDOU, K. 2000. Animal Bones and Animal Representations at Late Bronze Age Akrotiri. In: S. Sherratt (ed.), *Proceedings of the first International Symposium, The Wall Paintings of Thera*, vol. II, 709-735. Athens.
- VITRUVIUS 1960. *The Ten Books on Architecture*. Translated by Morris Hicky Morgan. New York.
- WALTER, H. & F. FELTEN. 1981. *Die vorgeschichtliche Stadt. Befestigungen. Häuser. Funde*. Alt-Ägina III.1. Mainz.
- WILD, E.M., W. GAUß, G. FORSTENPOINTNER, M. LINDBLOM, R. SMETANA, P. STEIER, U. THANHEISER & F. WENINGER 2010. 14C Dating of the Early to Late Bronze Age Stratigraphic Sequence of Aegina Kolonna, Greece. In: Breese, M.B.H., L.E. Rehn & C. Trautmann (eds), *Proceedings of the Eleventh International Conference on Accelerator Mass Spectrometry*, 1013-1021. Nuclear Instruments & Methods in Physics Research, Section B 268.
- ZEDER, M.A. & S.E. PILAAR 2010. Assessing the Reliability of Criteria Used to Identify Mandibles and Mandibular Teeth in Sheep, Ovis, and Goats, Capra. *Journal of Archaeological Science* 37, 225-242.
- ZEDER, M.A. & H.A. LAPHAM 2010. Assessing the Reliability of Criteria used to Identify Postcranial Bones in Sheep, Ovis, and Goats, Capra. *Journal of Archaeological Science* 37, doi:10.1016/j.jas.2010.06.032, 1-9.